

JOURNAL OF FUNCTIONAL ANALYSIS 12, 153-156 (1973)

 L^0 -Projective Topologies on Linear Spaces

H. MILLINGTON

*Mathematisches Institut der Universität Erlangen Nürnberg, 852 Erlangen
Bismarckstrasse 1^{1/2}, West Germany (BRD)**Communicated by I. Segal*

Received June 16, 1971

For any finite measure space (S, M, ν) let $L^0(S, M, \nu)$ denote the family of ν -equivalence classes of measurable functions on S , endowed with the topology of convergence in measure. For any topological vector space (X, \mathcal{G}) let $\lambda^0(X, \mathcal{G})$ denote the family of all continuous linear maps on X into (possibly different) spaces $L^0(S, M, \nu)$. If \mathcal{G} coincides with the coarsest topology on X making every map from $\lambda^0(X, \mathcal{G})$ continuous we call (X, \mathcal{G}) an L^0 -projective space. It is shown that L^p -spaces, $0 < p \leq 2$, and hence any projective limit of such spaces, are all L^0 -projective. It is further shown that l^p -spaces, $2 < p < \infty$, and the space c_0 , are not L^0 -projective.

The author particularly wishes to thank the referee for his helpful suggestions, especially for his shorter proof of Theorem 6, and the second observation of Theorem 1.

1. L^0 -PROJECTIVE SPACES

For any vector space X , and family \mathcal{L} of linear maps on X to possibly different topological vector spaces, $\mathcal{T}(\mathcal{L})$ is the coarsest topology on X making each map from \mathcal{L} continuous (the projective topology). For any topological vector space (X, \mathcal{G}) , X' is the topological dual.

DEFINITION. A topological vector space (X, \mathcal{G}) is L^0 -projective iff $\mathcal{T}(\lambda^0(X, \mathcal{G})) = \mathcal{G}$. (X, \mathcal{G}) is an L^0 -subspace iff there exists a linear homeomorphism from X onto a subspace of some $L^0(S, M, \nu)$.

In the terminology of L. Schwartz (10, 11), a space is L^0 -projective iff it has an 0-cotypical continuous map into some L^0 .^a

We make the following observations: Every $L^0(S, M, \nu)$ is a

topological vector space; hence, so also is $(X, \mathcal{T}(\lambda^0(X, \mathcal{G})))$ for any topological vector space (X, \mathcal{G}) (H. Schaefer, 9, p. 51). Clearly,

$$\mathcal{T}(\lambda^0(X, \mathcal{G})) \subset \mathcal{G}.$$

Also, by assigning to each $f \in X'$ the corresponding unit mass on X' , one checks that $\mathcal{T}(\lambda^0(X, \mathcal{G}))$ is finer than the weak topology on X .

In a straightforward manner one can prove also the following assertions.

THEOREM 1. *Every L^0 -subspace is L^0 -projective. A normed space is an L^0 -subspace iff it is an L^0 -projective space.*

THEOREM 2. *For any family \mathcal{L} of linear maps from a vector space X to L^0 -projective spaces, $\mathcal{T}(\mathcal{L})$ is L^0 -projective. In particular, any subspace, Cartesian product, or projective limit of L^0 -projective spaces is L^0 -projective.*

2. LINEAR STOCHASTIC PROCESSES

Without loss of generality we shall assume our measure spaces to be probability measure spaces. A linear map from a vector space X into an $L^0(S, M, \nu)$ is then called a *linear stochastic process*. We recall briefly certain well-known facts about such processes: cf. e.g. Gelfand and Vilenkin [5], Badrikian [1], Fernique [4], Cartier [3], Waldenfels [13].

For any linear process $A: X \rightarrow L^0(S, M, \nu)$, the *characteristic function* φ of A is defined by

$$\varphi(x) = \int e^{iA(x)(s)} d\nu(s), \quad x \in X.$$

Bochner's theorem has the following well-known generalization: A complex-valued function φ on a real vector space is the characteristic function of some linear process iff φ is positive-definite, $\varphi(0) = 1$, and φ is continuous on finite dimensional subspaces. It is also known that if φ is the characteristic function of a linear process A on a topological vector space (X, \mathcal{G}) then φ is continuous iff A is continuous. Hence,

THEOREM 3. *A topological vector space (X, \mathcal{G}) is L^0 -projective if there exists a positive-definite function φ on X such that*

$$x \rightarrow 0 \text{ in } X \quad \text{iff} \quad \varphi(x) \rightarrow 0.$$

For any measure space (S, M, ν) , finite or not, it is known that for $0 < p \leq 2$, $\varphi: f \in L^p(\nu) \rightarrow \exp(-\int |f|^p d\nu) \in \mathbb{C}$ is a positive definite-function. [2, Thm. 1, p. 232].

Thus, by Thms. 2, 3,

THEOREM 4. *Any L^p -space for $0 < p \leq 2$ is an L^0 -subspace. For any vector space X and family \mathcal{L} of linear maps from X into L^p -spaces, $0 < p \leq 2$, $\mathcal{T}(\mathcal{L})$ is an L^0 -projective topology on X .*

A space with a topology defined by a family of inner products (possibly with non-trivial kernels) is called *Hilbertian*. Such a space is necessarily a projective limit of L^2 -spaces [cf. 12, p. 115–116]. Also, every nuclear space is Hilbertian [12, p. 519]. Hence,

THEOREM 5. *Every Hilbertian space, in particular every nuclear space, is L^0 -projective.*

For each $1 \leq p < \infty$ let l^p be the Banach space of sequences (x_n) , $\sum_n |x_n|^p < \infty$, with the usual norm. Let c_0 be the space of all sequences (x_n) , $x_n \rightarrow 0$, with the sup norm. We shall now prove:

THEOREM 6. *c_0 , and the spaces l^p for $\infty > p > 2$, are not L^0 -projective. Any topological vector space with a subspace isomorphic to c_0 or an l^p -space, $\infty > p > 2$, is not L^0 -projective.*

Proof. Let $e_n: n = 0, 1, \dots$ be the canonical basis of c_0 , i.e. $e_{nm} = 1, n = m; e_{nm} = 0, n \neq m$. From Orlicz [8, Thm. 8(b), p. 300], or Kwapien [6], for any continuous linear process $A: c_0 \rightarrow L^0(S, M, \nu)$ we have that $\sum_n A(e_n)(s)^2 < \infty$ for ν -almost all $s \in S$. Thus $e_n \rightarrow 0$ in $\mathcal{T}(\lambda^0(c_0))$, but $e_n \not\rightarrow 0$ in the norm topology. Hence c_0 is not L^0 -projective.

Now, for any $(y_n) \in l^p$,

$$(x_n) \in c_0 \rightarrow (x_n y_n) \in l^p$$

is continuous. Hence, as a consequence of the Orlicz–Kwapien theorem, for any $A \in \lambda^0(l^p)$, $\sum_n A(e_n)^2 y_n^2$ converges ν -almost everywhere. Hence, if $p > 2$, since we must then have $(y_n^2) \in l^{p/2}$, it follows that for ν -almost $s \in S$, $(A(e_n)(s)^2) \in l^q$, where q is the index

conjugate to $p/2$, i.e. $(p/2)^{-1} + q^{-1} = 1$. Since $q < \infty$, then again $e_n \rightarrow 0$ in $\mathcal{T}(\lambda^0(l^p))$ but not in the norm topology.

The above theorem applies to suitable infinite-dimensional L^p -spaces, $\infty > p > 2$, and spaces $C(Y)$ of continuous functions.

With little extra effort the theorems given above all extend to the case of complex L^0 -spaces and complex vector spaces $X(7)$.

REFERENCES

1. A. BADRIKIAN, Séminaire sur les Fonctions Aléatoires Linéaires et les Mesures Cylindriques. Lecture Notes in Math. 139 (1970).
2. J. BRETAGNOLLE, D. DACUNHA-CASTELLE, AND J. L. KRIVINE, Lois stables et espaces L^p , *Ann. Inst. H. Poincaré Sec. B* 2 (1965-66), 231-259.
3. P. CARTIER, Processus aléatoires généralisés. Séminaire Bourbaki, 16e année: 1963/64, Fasc. 3, Exposé 272, Secrétariat mathématique, Paris 1964.
4. X. FERNIQUE, Processus linéaires, processus généralisés, *Ann. Inst. Fourier (Grenoble)* 17 (1967), 1-92.
5. I. M. GELFAND AND N. YA. VILENKIN, "Generalized Functions," Vol. 4, Academic Press, New York, 1965.
6. S. KWAPIEN, Complément au Théorème de Sazonov-Minlos, *Compt. Rend. Acad. Sci. Paris* 267 (1968), A698-700.
7. H. MILLINGTON, Cylinder measures over vector spaces, Thesis, U. British Columbia, 1971.
8. W. ORLICZ, On a class of asymptotically divergent sequences of functions, *Studia Math.* 12 (1951), 286-307.
9. H. H. SCHAEFER, Topological vector spaces, New York, MacMillan, 1966.
10. L. SCHWARTZ, Un théorème de dualité pour les applications radonifiantes, *Compt. Rend. Acad. Sci. Paris* 268 (9 Juin 1969), A1410-A1413.
11. L. SCHWARTZ, Un théorème de dualité pour les applications p -radonifiantes. Applications de ce théorème, in "Les Probabilités sur les Structures Algébriques," 319-326. C.N.R.S., Paris, 1970.
12. F. TREVES, "Topological Vector Spaces, Distributions and Kernels," Academic Press, 1966.
13. W. WALDENFELS, Positiv-definite Funktionen . . . *Studia Math.* 30 (1968), 153-161.